

## **How I treat Waldenstrom macroglobulinemia**

Steven P. Treon

Bing Center for Waldenstrom's Macroglobulinemia, Dana Farber Cancer Institute, and Department of Medicine, Harvard Medical School, Boston, MA 02115, USA.

### **Corresponding author:**

Steven P. Treon, M.D., M.A., Ph.D.

Bing Center for Waldenström's Macroglobulinemia

Dana Farber Cancer Institute

M548, 450 Brookline Avenue, Boston, MA 02115 USA

Tel: (617) 632-2681 Fax: (617) 632-4862

Email: [steven\\_treon@dfci.harvard.edu](mailto:steven_treon@dfci.harvard.edu)

**Short Title:** How I Treat Waldenstrom Macroglobulinemia.

**Keywords:** Waldenström macroglobulinemia, MYD88, CXCR4, rituximab, bortezomib, bendamustine, ibrutinib.

## Abstract

Waldenström's macroglobulinemia (WM) is a B-cell neoplasm manifested by the accumulation of clonal IgM secreting lymphoplasmacytic cells. MYD88 and CXCR4 WHIM-like somatic mutations are present in >90%, and 30-35% of WM patients, respectively, and impact disease presentation, treatment outcome, and/or overall survival. Familial predisposition is common in WM. Asymptomatic patients should be observed. Patients with disease related hemoglobin <10g/L, platelets <100x10<sup>9</sup>/L, bulky adenopathy and/or organomegaly, symptomatic hyperviscosity, peripheral neuropathy, amyloidosis, cryoglobulinemia, cold-agglutinin disease or transformed disease should be considered for therapy. Plasmapheresis should be used for patients with symptomatic hyperviscosity, and pre-rituximab for those with high serum IgM levels to pre-empt a symptomatic IgM flare. Treatment choice should take into account specific goals of therapy, necessity for rapid disease control, risk of treatment-related neuropathy, immunosuppression and secondary malignancies, and planning for future autologous stem cell transplantation. Frontline treatments include rituximab alone, or combined with alkylators (bendamustine, cyclophosphamide), proteasome-inhibitors (bortezomib, carfilzomib), nucleoside-analogues (fludarabine, cladribine) and ibrutinib. In the salvage setting, an alternative frontline regimen, ibrutinib, everolimus, or stem cell transplantation can be considered.

Investigational therapies under development for WM include agents that target MYD88, CXCR4, BCL2, and CD27/CD70 signaling, novel proteasome inhibitors, and chimeric antigen receptor modified T-cell therapy.

## Introduction

The cases presented in **Table 1** revealed Waldenström's macroglobulinemia (WM), a B-cell neoplasm resulting from the accumulation of clonal lymphoplasmacytic cells (LPC) secreting a monoclonal IgM protein.<sup>1,2</sup> WM patients can present with a wide array of symptoms and findings that impact diagnostic work-up and treatment. MYD88 and CXCR4 WHIM-like somatic mutations are present in >90%, and 30-35% of WM patients, respectively, but are absent or rare in other IgM secreting B-cell malignancies. Over half of individuals with IgM secreting monoclonal gammopathy of unknown significance (MGUS) harbor the MYD88<sup>L265P</sup> mutation, suggesting its role as an early oncogenic driver. Somatic CXCR4 mutations are similar to those found in the germline of WHIM (Warts, Hypogammaglobulinemia, Infections, Myelokathexis) syndrome, and nearly always are present in WM patients with MYD88<sup>L265P</sup>. Both nonsense (CXCR4<sup>WHIM/NS</sup>) and frameshift CXCR4 mutations (CXCR4<sup>WHIM/FS</sup>) occur in WM patients. MYD88 and CXCR4 mutations may impact disease presentation, treatment outcome, and/or survival.<sup>3-14</sup> Low tumor burden and serum IgM levels are associated with wild-type MYD88 (MYD88<sup>WT</sup>) disease, while MYD88<sup>L265P</sup> patients with CXCR4<sup>WHIM</sup> mutations have higher BM disease burden. High serum IgM levels including presentation with hyperviscosity crisis have been observed in

patients with CXCR4<sup>WHIM/FS</sup> tumor mutation status. Ibrutinib response (discussed below) is adversely impacted by MYD88<sup>WT</sup> and CXCR4<sup>WHIM</sup> status. In one study, overall survival was lower in patients with MYD88<sup>WT</sup> disease, but was unaffected by CXCR4 mutation status.

## Clinical Features

The clinical and laboratory findings for a large series of newly diagnosed patients who presented to the WM clinic at the Dana Farber Cancer Institute (a tertiary referral center) are depicted in **Table 2**. Splenomegaly and lymphadenopathy (as with Case 2) are uncommon at initial presentation (15-20%), though at later stages extra-medullary disease is more common (up to 60% of patients).<sup>16-18</sup> WM-related morbidity may be manifested by tumor cell infiltration, and/or by the physicochemical and immunological properties of the monoclonal IgM protein produced by WM cells (**Table 3**).<sup>19-</sup>

21

## Diagnostic Workup

### History taking

Familial predisposition is strong in WM.<sup>22-24</sup> Familial WM patients are younger, have a higher disease burden, and show lower response rates and progression-free survival (PFS) with non-proteasome inhibitor based therapy.<sup>25</sup> Hepatitis C exposure as a WM predisposition remains controversial though is associated with Type II cryoglobulinemia.<sup>26-28</sup> A thorough review of systems should be performed given a wide-range of WM-related morbidity that exists which can impact the diagnostic work-up and treatment undertaken (**Table 4**).

### **Laboratory Studies**

To establish the WM diagnosis, an IgM monoclonal protein and BM LPL infiltration must be present.<sup>1</sup> No minimum serum IgM or BM infiltration level is required to diagnose WM, since patients can be symptomatic and need treatment at low serum IgM (<1,000 mg/dL) levels or BM disease involvement.<sup>29</sup> Laboratory and clinical evaluations which can be performed in the initial work-up of all WM patients, and those with particular disease-related features are presented in **Table 4**.

## **Treatment Approaches to WM**

### **Management of the Asymptomatic WM Patient**

Patients with disease-related hemoglobin level (<10 g/dL), platelet count <100x10<sup>9</sup>/L, or symptomatic disease manifested by hyperviscosity, amyloidosis (particularly if there is heart involvement), cryoglobulinemia, cold-agglutininemia, extramedullary disease including central nervous system (Bing-Neel syndrome) involvement, moderate-severe, or advancing paraprotein-related peripheral neuropathy (PN), or disease transformation should be considered for therapy.<sup>30,31</sup> Treatment initiation should not be based on serum IgM levels *per se*, though at higher serum IgM levels (>6,000 mg/dL) empiric treatment may be appropriate to prevent hyperviscosity-related injury.

### **Management of the Symptomatic WM patient**

Treatment choice should take into account specific goals of therapy,

necessity for rapid disease control, risk of treatment-related neuropathy, immunosuppression, secondary malignancies, and potential for future autologous stem cell transplantation (ASCT). Heparin-related iron deficiency is common in WM patients.<sup>32</sup> In select patients with low disease burden, symptomatic anemia, and depressed iron saturation levels (i.e. <10-12%) unrelated to gastrointestinal bleeding, parenteral iron administration may be beneficial.<sup>32</sup> If unresponsive to parenteral iron, chemotherapy can then be initiated. Plasmapheresis should be considered for patients in need of immediate paraprotein control (discussed below), and pre-rituximab if serum IgM is  $\geq 4,000$  mg/dL to pre-empt a symptomatic IgM flare.<sup>30,33</sup> A suggested algorithm for the primary therapy of WM is provided in **Figure 1**. Continuous oral alkylator therapy (such as chlorambucil) and nucleoside-analogues should be avoided in ASCT candidates to prevent stem cell damage, as well as in younger patients (<65-70 years) given their association with secondary malignancies and/or disease transformation.<sup>30,33-37</sup>

### **Frontline Therapy of Waldenstrom's Macroglobulinemia**

For symptomatic, untreated WM patients not requiring immediate disease control as exemplified in Case 1, options include rituximab alone, or with

alkylators (bendamustine, cyclophosphamide), proteasome-inhibitors (bortezomib, carfilzomib), nucleoside-analogues (fludarabine, cladribine), and ibrutinib. Rituximab alone is well-suited for more indolent WM patients, i.e. those with mild-moderate symptomatic anemia or PN (as for Case 1), and those for whom more aggressive chemotherapy is inappropriate. Rituximab monotherapy should not be used in a patient in whom rapid responses are required, since response times can be slow. In addition, a rituximab-induced IgM flare (defined as  $\geq 25\%$  increase above baseline serum IgM level) occurs in 40-60% of WM patients and can provoke or accentuate IgM related morbidity including symptomatic hyperviscosity, PN, cold agglutininemia and cryoglobulinemia.<sup>38-42</sup> Rituximab monotherapy produces overall response rates (ORR) of 25-40% with standard (375 mg/m<sup>2</sup>/week for 4 weeks) therapy, and 40-60% with extended (375 mg/m<sup>2</sup>/week at weeks 1-4, 12-16) therapy with median PFS estimates of 16-29+ months.<sup>43-45</sup> Rituximab-related infusion reactions are common in WM patients, and administration of oral dexamethasone (10 mg) and famotidine (20 mg) the night before rituximab administration in addition to standard hypersensitivity prophylaxis may help reduce these events. Despite prophylaxis, progressive rituximab-intolerance is commonly encountered in WM, and ofatumumab may successfully be administered in place of rituximab.<sup>46,47</sup>

For untreated WM patients experiencing non-emergent paraprotein related morbidity, a proteasome-inhibitor containing regimen such as bortezomib, dexamethasone and rituximab (BDR) or carfilzomib, rituximab and dexamethasone (CaRD) is ideal.<sup>48-50</sup> Proteasome-inhibitor containing therapy can also be considered for myelosuppressed patients, though rarely bortezomib causes thrombocytopenia. In younger patients for whom avoidance of alkylator-based therapy is desired to reduce risk of secondary malignancies, proteasome-inhibitor based therapy (or ibrutinib discussed below) can be considered. Bortezomib is best avoided in patients with paraprotein-related neuropathy as in Case 1 given risk for treatment-related neuropathy. For BDR, weekly administration of bortezomib reduces but does not abrogate treatment-related neuropathy.<sup>49</sup> There is no data on subcutaneous bortezomib use in WM patients, but can be considered given the lower neuropathy risk observed with this method of administration in myeloma patients.<sup>51</sup> Regardless of route for administration, a close watch for bortezomib-related neuropathy should be undertaken, and drug held for grade  $\geq 2$  neuropathy. For weekly BDR, bortezomib can be dosed at 1.6 mg/m<sup>2</sup>/week with dexamethasone (20 mg/week) for 12-16 weeks, based on patient tolerance. Rituximab can be introduced if the baseline serum IgM is <4,000 mg/dL, or when such a level is achieved on weekly bortezomib and

dexamethasone alone. Plasmapheresis to lower the serum IgM to <4,000 mg/dL pre-rituximab can also be considered.<sup>30,33</sup> One single dose of rituximab (375 mg/m<sup>2</sup>/week) can be given every 3-4 weeks as part of the weekly BDR regimen. Serum IgM levels should be carefully monitored to detect a rituximab-related IgM flare. Maintenance can be considered for BDR responders (discussed below).

The CaRD regimen which is associated with a low incidence of treatment-related neuropathy can also be considered in non-emergent patients.<sup>50</sup> Serum IgA and IgG depletion is pronounced with CaRD which incorporates an extended maintenance schedule, and can contribute to recurring sinus and bronchial infections. A modified CaRD regimen with one rituximab infusion per induction cycle, or an abbreviated maintenance course may be considered to reduce serum IgA and IgG depletion.<sup>50</sup> Rituximab should only be introduced when the serum IgM is <4,000 mg/dL. Transient increases in amylase, lipase and bilirubin commonly occur with CaRD, and should be monitored. Carfilzomib should be avoided in patients at risk of cardiomyopathy. With either BDR or CaRD, herpes zoster prophylaxis with oral antiviral therapy (i.e. acyclovir 400 mg twice daily) for the duration of active therapy plus six months should be given, as well as famotidine (20

mg BID) during active therapy to decrease risk of gastrointestinal irritation.

For patients with bulky extramedullary disease, malignant pleural effusions or heavily impacted BM involvement, alkylator-based (bendamustine or cyclophosphamide) therapy can be considered. Benda-R showed longer (69 versus 29 months) PFS, and better tolerance versus CHOP-R in WM patients included in a randomized study by the German STiL group.<sup>52</sup> Bendamustine can be administered intravenously at 90 mg/m<sup>2</sup> on days 1, 2 every 4 weeks for 4-6 cycles, though 4 cycles is sufficient for most WM patients, and may reduce risk of prolonged or late myelosuppression. Bendamustine dose should be reduced in elderly patients (i.e. to 60-70 mg/m<sup>2</sup> on days 1, 2 every 4 weeks), and those with diminished renal function. Rituximab should be introduced when the serum IgM level is <4,000 mg/dL with Benda-R. A cyclophosphamide-based regimen such as rituximab, oral cyclophosphamide and dexamethasone (R-CD) can be considered as an alternative to Benda-R.<sup>30,33</sup> The ORR with R-CD or similar regimens are 80-90% with a median PFS of 3 years.<sup>53-55</sup> R-CHOP or R-CVP is not advantageous over R-CD or R-CP, and can contribute to more toxicities including treatment-related neuropathy.<sup>55</sup> Maintenance therapy following Benda-R or R-CD can be considered in responders (discussed below).

Nucleoside-analogues with rituximab and/or cyclophosphamide in untreated WM patients are associated with ORR of 85-95%, with a median PFS of 4 years.<sup>56-59</sup> It is unclear whether including cyclophosphamide to a nucleoside-analogue containing regimen (such as FCR) extends activity in WM, and may contribute to added toxicity. Because of immunosuppression, myelosuppression, impact on stem cell collection, and potential secondary malignancy risks including disease transformation, nucleoside-analogues should be cautiously used.<sup>34-37,56-59</sup> An exception may be for WM patients with Bing-Neel syndrome, for whom CNS responses can be obtained with fludarabine.<sup>60</sup> WM patients >70 years, and non-ASCT candidates may also be candidates for nucleoside-analogue therapy.<sup>31,34</sup> In elderly patients, fewer days of therapy per cycle, and fewer treatment cycles may be considered to minimize fludarabine-related toxicity.<sup>57</sup> Herpes zoster prophylaxis should be given for the duration of nucleoside-analogue treatment, and for at least one-year from end of active nucleoside-analogue therapy.<sup>57</sup> Cytokine support per ASCO guidelines should be considered with alkylator or nucleoside-analogue based therapies.<sup>61</sup>

Ibrutinib was recently approved by the U.S. Food and Drug Administration, and adopted into National Comprehensive Cancer Network

(NCCN) guidelines for symptomatic WM patients.<sup>30</sup> While data on the activity of ibrutinib in WM is based on previously treated patients (discussed below), the number of prior lines of therapy had no impact on its activity.<sup>14</sup> Moreover, less pre-treated WM patients showed better PFS, and less toxicity. However, patients with MYD88<sup>WT</sup> had considerably lower response rates and shorter PFS, while those with CXCR4 mutations showed slower initial response that improved with prolonged (>6 months) therapy.<sup>14</sup> The rapid time to IgM reduction (4 weeks), and absence of a rituximab-like IgM flare effect make the use of ibrutinib particularly attractive for patients with high IgM levels (in whom use of rituximab would be problematic), as well as those with IgM-related morbidity. There is no data on ibrutinib use in patients with Bing-Neel Syndrome, or bulky adenopathy (>5 cm), though reductions in extramedullary disease occurred in most previously treated patients on ibrutinib. The use of ibrutinib (if available) can be considered in symptomatic frontline patients, though alternative options should be considered for patients with bulky disease, and MYD88<sup>WT</sup> disease. Ibrutinib should be avoided in patients on warfarin, low platelet counts, and bleeding.<sup>14</sup> Patients with a prior history of arrhythmia may also experience tachyarrhythmias while on ibrutinib, and either close monitoring or alternative treatment options should be sought. To minimize bleeding risk associated with ibrutinib, any fish oil supplements should be discontinued, and drug

withheld for 3 to 7 days before and for 1 to 3 days after an anticipated invasive procedure.<sup>14</sup>

### **Treatment of the WM Patient Requiring Immediate Disease Control**

For WM patients requiring immediate paraprotein control (as in Case 2), plasmapheresis should initially be performed.<sup>30,33</sup> Typically 2-3 sessions of plasmapheresis reduce serum IgM levels by 30-60%.<sup>62,63</sup> Red blood cell transfusions (if required) should ideally follow plasmapheresis to not aggravate the viscosity load.<sup>63</sup> Blood-warmers should be used during plasmapheresis in patients with cryoglobulinemia or cold-agglutininemia to prevent cryoprecipitation and/or erythrocyte agglutination.

Treatment should be initiated as soon as possible after plasmapheresis, as serum IgM levels will return to baseline in 4-5 weeks.<sup>63</sup> Ibrutinib (if available) may be ideal in such a setting, given the rapid time to IgM reduction (median 4 and 8 weeks to at least a minor and major response, respectively). If ibrutinib is not available, modified BDR consisting of bortezomib at 1.3 mg/m<sup>2</sup> and dexamethasone 20 mg can be given twice weekly (on days 1, 4, 8, 11 every 21 days) for the first 1-2 cycles.<sup>49</sup> Once

the patient is stabilized, bortezomib (1.6 mg/m<sup>2</sup>) and dexamethasone (20 mg) can be given weekly, and up to 16 weeks (in total) of therapy can be considered. Rituximab (375 mg/m<sup>2</sup>/week) every 3-4 weeks can be integrated into BDR once the serum IgM is under 4,000 mg/dL. In some patients the serum IgM may remain unsafely high and continuance of twice-weekly bortezomib may be required. A close watch for bortezomib-related neuropathy should be maintained, and treatment held for grade  $\geq 2$  neuropathy. CaRD can also be considered which is associated with low treatment-related neuropathy risk.<sup>50</sup> As an alternative, bendamustine based therapy (as discussed above) can be considered.<sup>52</sup> The serum IgM should be carefully monitored for a few weeks after rituximab is initiated to monitor for an IgM flare. Maintenance therapy can be considered for responders and is discussed below.

### **Treatment of Paraprotein Related Peripheral Neuropathy**

For patients with mild, progressive IgM-related PN (such as Case 1), rituximab alone can be considered though its clinical impact remains in question.<sup>64-67</sup> Frequently, patients show a hematological response, without improvement in PN symptoms. The depth of response, and elapsed time

between manifestation of PN symptoms and therapy can be factors.<sup>68</sup> In patients with chronic, longstanding PN, the goal is to prevent symptomatic progression as myelin regeneration occurs very slowly. For patients with moderate to severe IgM-related PN, combination therapy with rituximab may also be effective.<sup>67,68</sup> Ibrutinib has shown symptomatic improvement in WM patients with IgM-related PN that progressed after rituximab, and is also reasonable.<sup>69</sup> In patients presenting with rapidly evolving IgM-related PN, interim plasmapheresis may be useful before instituting definitive chemotherapy. Steroid or IVIG use is of little value in the treatment of IgM-related PN.<sup>68</sup> Proteasome-inhibitor based therapy can be considered for amyloid or cryoglobulin related PN, using either weekly dosed bortezomib or carfilzomib to minimize risk of treatment-related neuropathy. As before, proteasome inhibitor therapy should be held for grade  $\geq 2$  treatment-related neuropathy. Cryoglobulin-related PN should only be treated if chronic symptoms are present, and not for transient, cold-induced episodic events.

### **Therapy in previously treated WM**

A suggested algorithm for the therapy of previously treated WM patients is provided in **Figure 2**. The overall response rate for ibrutinib in previously

treated WM patients was 91%, with 73% of these patients achieving a major response.<sup>14</sup> The 2-year estimate for progression-free and overall survival in this study was 69% and 95% respectively. Overall and major response rates were highest among patients with the MYD88<sup>L265P</sup>CXCR4<sup>WT</sup> genotype (100.0% and 91.2%), followed by MYD88<sup>L265P</sup>CXCR4<sup>WHIM</sup> (85.7% and 61.9%) and MYD88<sup>WT</sup>CXCR4<sup>WT</sup> (71.4% and 28.6%), respectively. Both major responders among the 7 patients with MYD88<sup>WT</sup>CXCR4<sup>WT</sup> were subsequently found to harbor other MYD88 mutations.<sup>70</sup>

Overall, ibrutinib was well tolerated, though neutropenia and thrombocytopenia were more common in heavily pre-treated patients. Serum IgA and IgG levels remained unchanged following ibrutinib, and treatment-related infections were infrequent. Patients should be maintained on ibrutinib until evidence of progression, or intolerance despite dose-reduction. Transient increases in serum IgM frequently occurred when ibrutinib was withheld, and did not signify treatment failure as IgM levels declined upon reinstatement of therapy.<sup>14</sup>

While ibrutinib represents an ideal treatment strategy for relapsed or refractory WM (as in Case 3), it is not available in many countries. The use

of an alternative first-line agent (as discussed above) can be considered if ibrutinib is not available. In patients for whom ASCT is seriously being considered, exposure to stem-cell damaging agents such as chlorambucil or nucleoside-analogues should be avoided, and a non-stem-cell toxic approach considered if stem cells have not been previously harvested.<sup>30,33</sup>

Everolimus has been investigated in WM. In previously treated WM patients, the overall and major response rates to everolimus were 73% and 50% respectively, with a median PFS of 21 months.<sup>69</sup> Grade  $\geq 3$  toxicities, particularly cytopenias, are common with everolimus. Pulmonary toxicity can also occur, and managed with steroids. Among untreated, symptomatic WM patients, the overall and major response rates for everolimus were 72% and 60%, respectively.<sup>71</sup> Discordance between serum IgM levels and BM disease response were commonly encountered in the frontline study which mandated serial bone marrow assessments. Oral ulcerations frequently occurred with everolimus, which could be prevented with an oral dexamethasone swish and spit solution. Because of its toxicity profile, and frequent toxicity-related treatment discontinuation, everolimus should be considered for third or beyond line of therapy in WM patients.

Stem cell transplantation (SCT) remains an option for salvage therapy in WM, particularly among younger patients who have had multiple relapses, or those patients with primary refractory disease. In an EBMTR study, the 5 year progression-free and overall survival were 40% and 69%, respectively, for patients with predominately multi-relapsed or refractory disease who received ASCT.<sup>72</sup> The non-relapse mortality at one-year was low (4%), and the presence of chemo-refractory disease and number of prior therapies impacted both progression-free and overall survival in this study. WM patients with amyloidosis may also benefit from ASCT.<sup>73</sup> The outcome of previously treated WM patients who received myeloablative and reduced-intensity allogeneic transplantation was also reported by the EBMTR.<sup>74</sup> Most patients (69%) in this series had chemotherapy-sensitive disease. The ORR was 76%, and the 5-year progression-free and overall survival rates were 56% and 62%, respectively. Among patients who received reduced-intensity allogeneic transplantation, similar progression-free and overall survival rates were observed (49% and 64%). Non-relapse mortality at 3 years was high, at 33% and 23% for myeloablative and reduced-intensity allogeneic transplantation, respectively. ASCT, and in very select cases, reduced-intensity allogeneic transplantation may be considered as appropriate salvage modalities, though the risks and benefits of these modalities should be carefully weighed against other available treatment options including

ibrutinib (if available).<sup>30,33</sup>

## **Maintenance Therapy in WM**

Maintenance rituximab therapy may be considered in WM patients responding to a rituximab-containing induction regimen.<sup>30,33</sup> In a large retrospective study, post-induction categorical responses improved in 42% of patients receiving maintenance rituximab versus 10% in those on watch-and-wait.<sup>75</sup> Progression-free and overall survival was longer for patients on maintenance rituximab as well. However, more infectious events (usually grade 1 or 2 sinusitis or bronchitis) and lower serum IgA and IgG levels were observed in patients on maintenance therapy. A prospective randomized study examining maintenance rituximab versus observation has been initiated by the German STiL group, and results awaited.<sup>76</sup>

The optimal dosing schedule for maintenance rituximab therapy in WM patients remains unclear. A single rituximab infusion (375 mg/m<sup>2</sup>) every 3 months for 2 years (as per Van Oers et al<sup>77</sup>) is reasonable over more frequent dosing to minimize rituximab-related serum IgA and IgG depletion.<sup>78</sup> Maintenance therapy should be discontinued in WM

patients with low serum IgG and/or IgA levels and recurring sinobronchial infections ( $\geq 3$ -4/year) that require antibiotic support, or who experienced a severe infection requiring hospitalization.

Combined maintenance therapy with rituximab and dexamethasone and either bortezomib or carfilzomib has been investigated in WM, and can be considered for WM patients who received BDR or CaRD induction, respectively. Serum IgA and IgG depletion may be more pronounced with proteasome-inhibitor containing maintenance regimens, and their levels should be closely monitored.<sup>48,50</sup> Shorter courses or longer intervals between cycles may help minimize serum IgA and IgG depletion, and can be considered in maintenance strategies that use BDR or CaRD. Neuropathy may also be potentiated with bortezomib containing maintenance regimens and should be carefully monitored, and treatment held for grade  $\geq 2$  treatment-related PN.<sup>48</sup>

## **Response Assessment**

Consensus response criteria have been developed as part of the International Workshops on WM.<sup>79</sup> A “major response” denotes a  $\geq 50\%$

decrease in serum IgM levels, and includes partial and complete responses, while the ORR includes minor responders. Durable clinical benefit has been shown in minor responders, while deeper categorical responses such as VGPR or CR are associated with longer PFS.<sup>80,81</sup> Response categories and criteria for progressive disease in WM are summarized in **Table 5**. Either the serum IgM or the IgM monoclonal spike may be used to track disease burden in WM patients.<sup>82</sup> However, gel migration patterns for the IgM paraprotein can vary and confound serial IgM monoclonal spike assessments in many cases.<sup>83</sup> In patients with cryoglobulinemia, serum samples should be obtained on warm baths (or equivalent) to minimize cryoprecipitation and serum IgM underestimation. Serum IgM levels can fluctuate with certain biological agents including rituximab, everolimus, bortezomib, and ibrutinib.<sup>14,38,39,70,84</sup> In circumstances when the serum IgM levels appear out of context with clinical impression, a BM biopsy should be considered.

## **Novel Agents for WM**

Investigational therapies under development for WM include agents that target MYD88, CXCR4, BCL2, and CD27/CD70 signaling, novel proteasome inhibitors, and chimeric antigen receptor (CAR) modified T-cell therapy. Idelalisib blocks PI3K $\delta$ , a growth promoting transcription factor that

is activated by MYD88<sup>L265P</sup>.<sup>85,86</sup> A 70% ORR was observed among ten previously treated WM patients evaluated in a larger study, and a prospective study in previously treated WM patients is forthcoming. IMO-8400 is an oligonucleotide that inhibits Toll-receptors 7, 8, and 9, and induces apoptosis in MYD88<sup>L265P</sup> WM cells.<sup>87</sup> A clinical trial examining IMO-8400 in WM has been initiated. IRAK1/IRAK4 kinases mediate MYD88<sup>L265P</sup> directed NFκB signaling, and their inhibition triggers apoptosis in MYD88<sup>L265P</sup> expressing malignant cells.<sup>7,88</sup> Moreover, combined BTK and IRAK1/4 inhibition induces synergistic killing of MYD88<sup>L265P</sup> malignant cells.<sup>88</sup> Compounds that inhibit IRAK signaling are under intense preclinical investigation for use in MYD88<sup>L265P</sup> diseases.<sup>89,90</sup> The anti-apoptotic factor BCL-2 is over-expressed in WM cells.<sup>91,92</sup> The BCL-2 inhibitor ABT-199 induces apoptosis, and shows at least additive anti-apoptotic activity against WM cells co-treated with either ibrutinib or idelalisib.<sup>93</sup> In a prospective clinical study, 3 of 4 WM previously treated WM patients responded, that included one CR.<sup>94</sup>

Oprozomib is an oral epoxyketone proteasome-inhibitor that is an analog of carfilzomib.<sup>95</sup> An ORR of 59% was observed in a phase 2 study in previously treated WM patients.<sup>96</sup> Clinical trials examining the oral proteasome-inhibitor ixazomib have also been initiated in combination with dexamethasone

and rituximab in symptomatic untreated, as previously treated WM patients.<sup>97,98</sup> Soluble CD27 (sCD27) is released by WM cells, and stimulates growth promoting ligands through CD70.<sup>99</sup> CD70 is also expressed on WM cells.<sup>100</sup> A clinical trial examining a novel human anti-CD70 (ArgenX-70) antibody that mediates ADCC and blocks sCD27-CD70 signaling is being initiated in previously treated WM patients.<sup>101</sup> In pre-clinical studies, CXCR4 antagonists blocked SDF-1a rescue of apoptosis mediated by ibrutinib, idelalisib, and other therapeutics.<sup>10,102,103</sup> A trial examining ibrutinib with the anti-CXCR4 antibody ulocuplumab (BMS-936564) is being planned in WM patients given encouraging safety and efficacy data in myeloma patients, as well as *in vivo* results with ulocuplumab in mice engrafted with CXCR4<sup>WHIM</sup> mutated WM cells.<sup>10,104</sup> Chimeric antigen receptor (CAR) modified T-cell therapy utilizing a second generation CAR derived from a CD19-directed antibody fused to the  $\zeta$  chain of CD3, and the intracellular signaling domain of CD28 (19-28z) has shown robust preclinical activity against WM cells.<sup>105</sup> A clinical trial for relapsed or refractory WM patients using 19-28z CAR modified autologous T-cells has been initiated.

## Case Management

**Case 1.** Given the non-urgent nature of this WM patient's presentation, rituximab was started for progressive, symptomatic IgM-related neuropathy. The patient did encounter an IgM flare, and worsening neuropathic symptoms. He underwent plasmapheresis and was transitioned to CDR, and achieved a PR with near resolution of his neuropathic symptoms. He completed 2-years of maintenance rituximab therapy, and remains in a partial response, without peripheral neuropathy, five-years later. Under similar circumstances today, ibrutinib (if available) could be considered as an appropriate initial or second line intervention.

**Case 2.** This 42-year old WM patient presented with symptomatic hyperviscosity. The patient also had bulky adenopathy and splenomegaly at the time of presentation. He underwent emergent plasmapheresis, and then received blood transfusions so as to not aggravate whole blood viscosity levels. Given his young age, and need for rapid paraprotein reduction, bortezomib and dexamethasone (without rituximab to avoid an IgM-flare) was started. Serum IgM levels were closely monitored post-pheresis, and he received additional plasma exchanges when serum IgM exceeded 5,000

mg/dL. After three cycles of bortezomib and dexamethasone no response occurred, and he was started on bendamustine. The serum IgM level decreased to 3,400 mg/dL following 2 cycles of bendamustine, and no further plasmapheresis was required. Rituximab was then added to bendamustine for 4 additional cycles, with continued close monitoring of serum IgM levels. Maintenance rituximab therapy followed for 2 additional years, and he achieved a VGPR with complete resolution of all extra-medullary disease. Two-years after treatment, he remains without progression, though has required IVIG replacement for recurring sino-bronchial infections associated with treatment related serum IgA and IgG depletion. Under similar circumstances today, ibrutinib (if available), would represent a very reasonable option to use either as initial or subsequent therapy. The IgG and IgA sparing effects associated with ibrutinib may have also prevented the need for IVIG replacement in this patient.

**Case 3.** This case illustrates a young WM patient refractory to multiple agents. Nucleoside analogues, alemtuzumab and allogeneic stem cell transplantation were considered, though the patient declined these options because of toxicity concerns. A BM biopsy confirmed persistent WM disease with a MYD88<sup>L265P</sup>CXCR4<sup>WT</sup> genotype. He received ibrutinib on a clinical trial and attained a partial response. Post- therapy, his hematocrit normalized

and he remains in a partial response 2.5 years later.

## **Acknowledgements**

This work was supported by Peter Bing M.D., the Linda and Edward Nelson Endowment for Studies into Waldenstrom's Macroglobulinemia, the Kerry Robertson Fund for Waldenstrom's Research, the Bauman Family Foundation, and the International Waldenstrom's Macroglobulinemia Foundation.

## **Conflicts of Interest**

Dr. Treon has disclosed active research funding, consulting fees, and/or speaking honoraria from Janssen Pharmaceuticals Inc., Onyx Inc., Pharmacyclics Inc., and Gilead Pharmaceuticals Inc.

## References

1. Owen RG, Treon SP, Al-Katib A, et al. Clinicopathological definition of Waldenström's macroglobulinemia: Consensus Panel Recommendations from the Second International Workshop on Waldenström's macroglobulinemia. *Semin Oncol* 2003; 30:110–15.
2. World Health Organization Classification of Tumours of Haematopoietic and Lymphoid Tissues. Swerdlow, SH, Campo, E, Harris, NL, et al. (Eds), IARC Press, Lyon 2008.
3. Treon SP, Xu L, Yang G, et al: MYD88 L265P somatic mutation in Waldenstrom's macroglobulinemia. *N Engl J Med* 2012; 367:826-33.
4. Xu L, Hunter Z, Yang G, et al. MYD88 L265P in Waldenstrom macroglobulinemia, immunoglobulin M monoclonal gammopathy, and other B-cell lymphoproliferative disorders using conventional and quantitative allele-specific polymerase chain reaction. *Blood*. 2013; 121(11):2051-8.
5. Varettoni M, Arcaini L, Zibellini S, et al. Prevalence and clinical significance of the MYD88 L265P somatic mutation in Waldenstrom macroglobulinemia, and related lymphoid neoplasms. *Blood*. 2013; 121: 2522-8.
6. Jiménez C, Sebastián E, Del Carmen Chillón M, et al. MYD88 L265P is a marker highly characteristic of, but not restricted to, Waldenström's

- macroglobulinemia. *Leukemia* 2013; 27(8):1722-8.
7. Poulain S, Roumier C, Decambron A, et al. MYD88 L265P mutation in Waldenstrom's macroglobulinemia. *Blood* 2013; 121(22): 4504-11.
  8. Ansell SM, Hodge LS, Secreto FJ, et al. Activation of TAK1 by MYD88 L265P drives malignant B-cell growth in Non-Hodgkin lymphoma. *Blood Cancer J.* 2014 Feb 14;4:e183. doi: 10.1038/bcj.2014.4.
  9. Hunter ZR, Xu L, Yang G, Zhou Y, Liu X, Cao Y, et al. The genomic landscape of Waldenstrom's Macroglobulinemia is characterized by highly recurring MYD88 and WHIM-like CXCR4 mutations, and small somatic deletions associated with B-cell lymphomagenesis. *Blood* 2014; 123(11):1637-46.
  10. Roccaro A, Sacco A, Jiminez C, et al. C1013G/CXCR4 acts as a driver mutation of tumor progression and modulator of drug resistance in lymphoplasmacytic lymphoma. *Blood.* 2014; 123(26):4120-31.
  11. Schmidt J, Federmann B, Schindler N, et al. MYD88 L265P and CXCR4 mutations in lymphoplasmacytic lymphoma identify cases with high disease activity. *Br J Haematol* 2015 Mar 29. doi: 10.1111/bjh.13361. [Epub ahead of print]
  12. Poulain S, Roumier C, Doye E, et al. Genomic Landscape of CXCR4 Mutations in Waldenstrom's Macroglobulinemia. *Blood* 2014 124:1627.
  13. Treon SP, Cao Y, Xu L, et al. Somatic mutations in MYD88 and CXCR4 are determinants of clinical presentation and overall survival in

- Waldenstrom macroglobulinemia. *Blood*. 2014;123(18):2791-6.
14. Treon SP, Tripsas CK, Meid K, et al. Ibrutinib in previously treated Waldenstrom's Macroglobulinemia. *N Engl J Med*. 2015; 372(15):1430-40.
  15. Treon SP, Hunter ZR, Matous J, et al. Multicenter clinical trial of bortezomib in relapsed/refractory Waldenstrom's macroglobulinemia: results of WMCTG Trial 03-248. *Clin Cancer Res*. 2007; 13(11):3320-5.
  16. Ghobrial IM, Gertz M, LaPlant B, et al. Phase II Trial of the Oral Mammalian Target of Rapamycin Inhibitor Everolimus in Relapsed or Refractory Waldenström Macroglobulinemia. *J Clin Oncol*. 2010; 28(8): 1408–1414.
  17. Tedeschi A, Benevolo G, Varettoni M, et al. Fludarabine plus cyclophosphamide and rituximab in Waldenstrom macroglobulinemia: an effective but myelosuppressive regimen to be offered to patients with advanced disease. *Cancer*. 2012; 118(2):434-43.
  18. Merlini G, Farhangi M, Osserman EF. Monoclonal immunoglobulins with antibody activity in myeloma, macroglobulinemia and related plasma cell dyscrasias. *Semin Oncol* 1986; 13:350–65.
  19. Farhangi M, Merlini G. The clinical implications of monoclonal immunoglobulins. *Semin Oncol* 1986; 13:366–79.
  20. Marmont AM, Merlini G. Monoclonal autoimmunity in hematology. *Haematologica* 1991; 76:449–59.

21. Treon SP, Hunter ZR, Aggarwal A, et al. Characterization of familial Waldenstrom's Macroglobulinemia. *Ann Oncol* 2006; 17: 488-94.
22. McMaster ML, Csako G, Giambarresi TR, et al. Long-term evaluation of three multiple-case Waldenstrom macroglobulinemia families. *Clin Cancer Res* 2007; 13: 5063-9.
23. Kristinsson SY, Bjorkholm M, Goldin LR, et al. Risk of lymphoproliferative disorders among first-degree relatives of lymphoplasmacytic lymphoma/Waldenstrom macroglobulinemia patients: a population-based study in Sweden. *Blood* 2008; 112: 3052-6.
24. Treon SP, Tripsas C, Hanzis C, et al. Familial disease predisposition impacts treatment outcome in patients with Waldenström macroglobulinemia. *Clin Lymphoma Myeloma Leuk.* 2012;12(6):433-7.
25. Santini GF, Crovatto M, Modolo ML, et al. Waldenström macroglobulinemia: a role of HCV infection? *Blood* 1993; 82:2932.
26. Silvestri F, Barillari G, Fanin R, et al. Risk of hepatitis C virus infection, Waldenström's macroglobulinemia, and monoclonal gammopathies. *Blood* 1996; 88:1125-6.
27. Leleu X, O'Connor K, Ho A, et al. Hepatitis C Viral Infection Is Not Associated with Waldenstrom's Macroglobulinemia. *Am J Hematol* 2006. *Am J Hematol* 2007; 82: 83-4.
28. Treon SP. How I treat Waldenstrom Macroglobulinemia. *Blood.* 2009; 114:2375-85.

29. Kyle RA, Treon SP, Alexanian R, et al. Prognostic markers and criteria to initiate therapy in Waldenstrom's Macroglobulinemia: Consensus Panel Recommendations from the Second International Workshop on Waldenstrom's Macroglobulinemia. *Semin Oncol* 2003; 30: 116-120.
30. Anderson KC, Alsina M, Bensinger W, et al. Waldenström's macroglobulinemia/lymphoplasmacytic lymphoma, version 1.2013. *J Natl Compr Canc Netw*. 2013 Jan 1;11(1):11-7.32.
31. Ciccarelli BT, Patterson CJ, Hunter ZR, et al. Hepcidin is produced by lymphoplasmacytic cells and is associated with anemia in Waldenström's Macroglobulinemia. *Clin Lymphoma Myeloma Leuk*. 2011; 11(1):160-3.
32. Treon SP, Tripsas C, Ciccarelli BT, Manning RJ, Patterson CJ, Sheehy P, Hunter ZR. Patients with Waldenstrom Macroglobulinemia Commonly Present With Iron Deficiency and Those With Severely Depressed Transferrin Saturation Levels Show Response to Parenteral Iron Administration. *Clin Lymphoma Myeloma Leuk*. 2013; 13(2):241-3.
33. Dimopoulos MA, Kastiris E, Owen RG et al. Treatment recommendations for patients with Waldenström macroglobulinemia (WM) and related disorders: IWWM-7 consensus. *Blood*. 2014; 124(9):1404-11.
34. Thomas S, Hosing C, Delasalle KB, et al. Success rates of autologous stem cell collection in patients with Waldenstrom's macroglobulinemia. *Proc 5th International Workshop on Waldenstrom's macroglobulinemia 2008 (Supplemental Abstract)*.

35. Leleu XP, Manning R, Soumerai JD, et al. Increased incidence of transformation and myelodysplasia/acute leukemia in patients with Waldenström macroglobulinemia treated with nucleoside analogs. *J. Clin. Oncol.* 2009; 27: 250-5.
36. Leleu X, Tamburini J, Roccaro A, et al. Balancing risk versus benefit in the treatment of Waldenström's macroglobulinemia patients with nucleoside-analogue based therapy. *Clin. Lymph. Myeloma.* 2009; 9:71-73.
37. Rakkhit R, Delasalle KB, Gavino MB, et al. Incidence of transformation to large cell lymphoma and to second malignancies in symptomatic patients with Waldenström's macroglobulinemia (WM) treated with cladribine (2-CdA) combination induction. *Blood.* 2008; 112: Abstract 3065.
38. Treon SP, Branagan AR, Hunter ZR, et al. Paradoxical increases in serum IgM and viscosity levels following rituximab in Waldenström's macroglobulinemia. *Ann Oncol* 2004; 15:1481-3.
39. Ghobrial IM, Fonseca R, Greipp PR, et al: Initial immunoglobulin M "flare" after rituximab therapy in patients with Waldenström Macroglobulinemia: An Eastern Cooperative Oncology Group Study. *Cancer* 2004; 101:2593-8.
40. Ghobrial IM, Uslan DZ, Call TG, et al. Initial increase in the cryoglobulin level after rituximab therapy for type II cryoglobulinemia secondary to Waldenström macroglobulinemia does not indicate failure of

- response. *Am J Hematol* 2004; 77: 329-30.
41. Noronha V, Fynan TM, Duffy T. Flare in neuropathy following rituximab therapy for Waldenstrom's macroglobulinemia. *J Clin Oncol* 2006; 24: E3.
  42. Broglio L, Lauria G. Worsening after rituximab treatment in anti-MAG neuropathy. *Muscle Nerve* 2005; 32:378-9.
  43. Gertz MA, Rue M, Blood E, et al: Multicenter phase 2 trial of rituximab for Waldenstrom macroglobulinemia (WM): An Eastern Cooperative Oncology Group Study (E3A98) *Leuk. Lymphoma*. 45:2047-2055, 2004.
  44. Dimopoulos MA, Zervas C, Zomas A, et al: Treatment of Waldenstrom's macroglobulinemia with rituximab. *J Clin Oncol* 2002; 20:2327-33.
  45. Treon SP, Emmanouilides C, Kimby E, et al. Extended rituximab therapy in Waldenström's Macroglobulinemia. *Ann Oncol* 2005; 16:132-8.
  46. Furman RR, Eradat H, Switzky JC, et al. A Phase II Trial of Ofatumumab In Subjects with Waldenstrom's Macroglobulinemia. *Blood* 2011; 118: Abstract 3701.
  47. Kanan S, Meid K, Treon SP, et al. Clinical Characteristics of Rituximab Intolerance in Patients with Waldenstrom's Macroglobulinemia. *Proc. Am. Soc. Hematology* 2014; Abstract 2610.
  48. Treon SP, Ioakimidis L, Soumerai JD, et al. Primary therapy of Waldenstrom's Macroglobulinemia with Bortezomib, Dexamethasone and

- Rituximab: Results of WMCTG Clinical Trial 05-180. *J. Clin. Oncol.* 2009; 27:3830-5.
49. Dimopoulos MA, García-Sanz R, Gavriatopoulou M, et al. Primary therapy of Waldenstrom macroglobulinemia (WM) with weekly bortezomib, low-dose dexamethasone, and rituximab (BDR): long-term results of a phase 2 study of the European Myeloma Network (EMN). *Blood.* 2013; 122(19):3276-82.
50. Treon SP, Tripsas CK, Meid K, et al. Carfilzomib, rituximab, and dexamethasone (CaRD) treatment offers a neuropathy-sparing approach for treating Waldenström's Macroglobulinemia. *Blood.* 2014; 124(4):503-10.
51. Moreau P, Pylypenko H, Grosicki S, Karamanesht I, et al. Subcutaneous versus intravenous administration of bortezomib in patients with relapsed multiple myeloma: a randomised, phase 3, non-inferiority study. *Lancet Oncol.* 2011; 12(5):431-40.
52. Rummel M, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet.* 2013; 381(9873):1203-10.
53. Dimopoulos MA, Anagnostopoulos A, Kyrtonis MC, et al. Primary treatment of Waldenstrom's macroglobulinemia with

- Dexamethasone, Rituximab and Cyclophosphamide. *J Clin Oncol* 2007; 25:3344-9.
54. Buske C, Hoster E, Dreyling MH, et al. The addition of rituximab to front-line therapy with CHOP (R-CHOP) results in a higher response rate and longer time to treatment failure in patients with lymphoplasmacytic lymphoma: results of a randomized trial of the German Low-Grade Lymphoma Study Group (GLSG). *Leukemia* 2009; 23: 153-61.
55. Ioakimidis L, Patterson CJ, Hunter ZR, Soumerai JD, Manning RJ, Sheehy P, Treon SP. Comparative outcomes following CP-R, CVP-R and CHOP-R in Waldenstrom's Macroglobulinemia. *Clin Lymphoma Myeloma* 2009; 9:62-6.
56. Weber DM, Dimopoulos MA, Delasalle K, et al. 2-Chlorodeoxyadenosine alone and in combination for previously untreated Waldenstrom's macroglobulinemia. *Semin Oncol*. 2003 Apr;30(2):243-7.
57. Treon SP, Branagan AR, Ioakimidis L, et al. Long term outcomes to Fludarabine and Rituximab in Waldenström's macroglobulinemia. *Blood* 2009; 113(16): 3673-8.
58. Laszlo D, Andreola G, Rigacci L, et al. Rituximab and Subcutaneous 2-Chloro-2'-Deoxyadenosine Combination Treatment for Patients With Waldenström Macroglobulinemia: Clinical and Biologic Results of a Phase II Multicenter Study. *J. Clin. Oncol*. 2010; 28(13):2233-8.
59. Tedeschi A, Benevolo G, Varettoni M, et al. Fludarabine plus

- cyclophosphamide and rituximab in Waldenstrom macroglobulinemia: an effective but myelosuppressive regimen to be offered to patients with advanced disease. *Cancer*. 2012; 118(2):434-43.
60. Vos J MI, Kersten MJ, Kraan W, et al. Bing Neel Syndrome with good response to Low Dose oral Fludarabine in Four (4) Consecutive Patients. 8th International Workshop on Waldenstrom's Macroglobulinemia. August 14-16, 2014. Abstract W25.
61. Smith TJ, Khatcheressian J, Lyman GH, et al. 2006 update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline. *J Clin Oncol*. 2006; 24(19):3187-205.
62. Stone MJ, Bogen SA. Evidence-based focused review of management of hyperviscosity syndrome. *Blood*. 2012; 119(10):2205-8.
63. Menke MN, Treon SP. Hyperviscosity Syndrome. In: *Clinical Malignant Hematology*, pgs. 937-41, Sekeres, Kalaycio, Bolwell eds., McGraw Hill Publishing, New York, 2007.
64. Renaud S, Gregor M, Fuhr P, et al. Rituximab in the treatment of polyneuropathy associated with anti-MAG antibodies. *Muscle Nerve*. 2003; 27(5):611-5.
65. Léger JM, Viala K, Nicolas G, et al. Placebo-controlled trial of rituximab in IgM anti-myelin-associated glycoprotein neuropathy. *Neurology*. 2013; 80(24):2217-25.

66. Lunn MP, Nobile-Orazio E. Immunotherapy for IgM anti-myelin-associated glycoprotein paraprotein-associated peripheral neuropathies. *Cochrane Database Syst Rev.* 2012; 5:CD002827. doi: 10.1002/14651858.CD002827.pub3.
67. Hospital MA, Viala K, Dragomir S, et al. Immunotherapy-based regimen in anti-MAG neuropathy: results in 45 patients. *Haematologica.* 2013;98(12):e155-7. doi: 10.3324/haematol.2013.087213.
68. Treon SP, Hanzis C, Ioakimidis L, et al. Clinical characteristics and treatment outcome of disease-related peripheral neuropathy in Waldenstrom's macroglobulinemia (WM). *J Clin Oncol.* 2010; 28:15s (Abstract 8114).
69. Ghobrial IM, Witzig TE, Gertz M, et al. Long-term results of the phase II trial of the oral mTOR inhibitor everolimus (RAD001) in relapsed or refractory Waldenstrom Macroglobulinemia. *Am J Hematol.* 2014; 89(3):237-42.
70. Treon SP, Xu L, Hunter ZR. MYD88 mutations and ibrutinib responses in Waldenstrom's Macroglobulinemia. *N Engl J Med.* 2015; Letter in press.
71. Treon SP, Tripsas CK, Meid K, et al. Prospective, multicenter study of the MTOR inhibitor everolimus (RAD001) as primary therapy in Waldenstrom's Macroglobulinemia. *Blood* 2013; 122:1822.
72. Kyriakou C, Canals C, Sibon D, et al. High-dose therapy and autologous stem-cell transplantation in Waldenstrom macroglobulinemia:

- the Lymphoma Working Party of the European Group for Blood and Marrow Transplantation. *J Clin Oncol* 2010; 28:2227-32.
73. Gertz MA, Hayman SR, Buadi FK. Transplantation for IgM amyloidosis and IgM myeloma. *Clin Lymphoma Myeloma* 2009; 9:77-79.
74. Kyriakou C, Canals C, Cornelissen JJ, et al. Allogeneic stem-cell transplantation in patients with Waldenström macroglobulinemia: report from the Lymphoma Working Party of the European Group for Blood and Marrow Transplantation. *J Clin Oncol*. 2010; 28:4926-34.
75. Treon SP, Hanzis, C, Manning, RJ, et al. Maintenance Rituximab is associated with improved clinical outcome in Rituximab naïve patients with Waldenstrom's Macroglobulinemia who respond to a Rituximab containing regimen. *Br J Haematol* 2011; 154:357-62.
76. Rummel MJ, Lerchenmüller C, Greil R, et al. Bendamustine-Rituximab Induction Followed by Observation or Rituximab Maintenance for Newly Diagnosed Patients with Waldenström's Macroglobulinemia: Results From a Prospective, Randomized, Multicenter Study (StiL NHL 7-2008). *Blood* 2012; 120: 2739.
77. van Oers MH, Klasa R, Marcus RE, et al. Rituximab maintenance improves clinical outcome of relapsed/resistant follicular non-Hodgkin lymphoma in patients both with and without rituximab during induction: results of a prospective randomized phase 3 intergroup trial. *Blood* 2006; 108:3295-301.

78. Hunter ZR, Manning RJ, Hanzis C, et al. IgA and IgG hypogammaglobulinemia in Waldenström's macroglobulinemia. *Haematologica*. 2010; 95(3):470-5.
79. Owen RG, Kyle RA, Stone MJ, et al. Response Assessment in Waldenstrom Macroglobulinemia. *Brit. J. Haematol*. 2013; 160(2):171-6.
80. Gertz MA, Abonour R, Heffner LT, et al. Clinical value of minor responses after 4 doses of rituximab in Waldenström macroglobulinaemia: a follow-up of the Eastern Cooperative Oncology Group E3A98 trial. *Brit. J. Haematol*. 2009; 147(5):677-80.
81. Treon SP, Yang G, Hanzis C, et al. Attainment of complete/very good partial response following rituximab-based therapy is an important determinant to progression-free survival, and is impacted by polymorphisms in FCGR3A in Waldenstrom macroglobulinaemia. *Brit. J. Haematol*. 2011; 154(2): 223-8.
82. Tripsas CK, Patterson CJ, Uljon SN, et al. Comparative response assessment by serum immunoglobulin M M-protein and total serum immunoglobulin M after treatment of patients with Waldenström macroglobulinemia. *Clin Lymphoma Myeloma Leuk*. 2013; 13(2):250-2.
83. Uljon SN, Treon SP, Tripsas CK, Lindeman NI. Challenges with serum protein electrophoresis in assessing progression and clinical response in patients with Waldenström macroglobulinemia. *Clin Lymphoma Myeloma Leuk*. 2013; 13(2):247-9.

84. Strauss SJ, Maharaj L, Hoare S, et al. Bortezomib therapy in patients with relapsed or refractory lymphoma: Potential correlation of in vitro sensitivity and tumor necrosis factor alpha response with clinical activity. *J Clin Oncol* 2006; 24: 2105-12.
85. Yang G, Liu X, Zhou Y, et al. PI3K/AKT pathway is activated by MYD88 L265P and use of PI3K-delta inhibitors induces robust tumor cell killing in Waldenstrom's Macroglobulinemia. *Blood* 2013; 121: Abstract 4255.
86. Gopal AK, Kahl BS, de Vos S, Wagner-Johnston ND, Schuster SJ, Jurczak WJ et al. PI3K $\delta$  inhibition by idelalisib in patients with relapsed indolent lymphoma. *N Engl J Med*. 2014; 370(11):1008-18.
87. Brenner L, Arbeit RD, Sullivan T. IMO-8400, an Antagonist of Toll-like Receptors 7, 8, and 9, in Development for Genetically Defined B-Cell Lymphomas: Safety and Activity in Phase 1 and Phase 2 Clinical Trials. *Proc. American Society for Hematol*. 2014; Abstract 3101.
88. Yang G, Zhou Y, Liu X, Xu L, Cao Y, Manning RJ, et al. A mutation in MYD88 (L265P) supports the survival of lymphoplasmacytic cells by activation of Bruton tyrosine kinase in Waldenstrom macroglobulinemia. *Blood* 2013; 122(7):1222-32.
89. Lim KH, Romero DL, Chaudhary D, et al. IRAK4 Kinase As A Novel Therapeutic Target in the ABC Subtype of Diffuse Large B Cell Lymphoma; *Proc. Am. Soc. Hematol*. 2012; Abstract 62.
90. Burhlage S. Kinome targets and inhibitors. 8th International

Workshop on Waldenstrom's Macroglobulinemia. August 14-16, 2014.  
Abstract 27.

91. Chng WJ, Schop RF, Price-Troska T, et al. Gene-expression profiling of Waldenström macroglobulinemia reveals a phenotype more similar to chronic lymphocytic leukemia than multiple myeloma. *Blood* 2006; 108:2755–63.
92. Hatjiharissi, E, Mitsiades CS, Ciccarelli B, et al. Comprehensive Molecular Characterization of malignant and microenvironmental cells in Waldenstrom's Macroglobulinemia by Gene Expression Profiling. *Blood* 2007; 110:933a.
93. Cao Y, Yang G, Hunter ZR, et al. The BCL2 antagonist ABT-199 triggers apoptosis, and augments ibrutinib and idelalisib mediated cytotoxicity in CXCR4Wild-type and CXCR4WHIM mutated Waldenstrom macroglobulinaemia cells. *Brit. J. Hematol.* 2014; manuscript accepted.
94. Davids MS, Seymour JF, Gerecitano JF, et al. Phase I study of ABT-199 (GDC-0199) in patients with relapsed/refractory non-Hodgkin lymphoma: Responses observed in diffuse large B-cell (DLBCL) and follicular lymphoma at higher cohort doses. *J. Clin. Oncol.* 2015; 32(5s): Abstract 8522.
95. Hurchla MA, Garcia-Gomez A, Hornick MC, et al. The epoxyketone-based proteasome inhibitors carfilzomib and orally bioavailable oprozomib have anti-resorptive and bone-anabolic activity in addition to anti-myeloma

- effects. *Leukemia*. 2013; 27(2):430-40.
96. Siegel DS, Kaufman JL, Raje NS, et al. Updated Results from a Multicenter, Open-Label, Dose-Escalation Phase 1b/2 Study of Single-Agent Oprozomib in Patients with Waldenström Macroglobulinemia (WM). *Proc. American Society for Hematol*. 2014; Abstract 1785.
97. Chauhan D, Tian Z, Zhou B, et al. In vitro and in vivo selective antitumor activity of a novel orally bioavailable proteasome inhibitor MLN9708 against multiple myeloma cells. *Clin Cancer Res* 2011; 17(16):5311-5321.
98. Richardson PG, Baz R, Wang M. Phase 1 study of twice-weekly ixazomib, an oral proteasome inhibitor, in relapsed/refractory multiple myeloma patients. *Blood*. 2014; 124(7):1038-46.
99. Sahota SS, Forconi F, Ottensmeier CH, et al. Typical Waldenström macroglobulinemia is derived from a B-cell arrested after cessation of somatic mutation but prior to isotype switch events. *Blood* 2002; 100:1505-7.
100. Ho AW, Hatjiharissi E, Branagan AR, et al. CD27-CD70 interactions in the pathogenesis of Waldenström's Macroglobulinemia. *Blood* 2008; 112(12):4683-9.
101. Thibault A, de Haard H, Meid K, et al. A Phase 1/2 study of ARGX-110, a monoclonal antibody targeting CD70, a receptor involved in sCD27 signaling in patients with relapsed or refractory Waldenström's

- macroglobulinemia. 8th International Workshop on Waldenstrom's Macroglobulinemia. August 14-16, 2014. Abstract W38.
102. Cao Y, Hunter ZR, Liu X, et al. The WHIM-like CXCR4S338X somatic mutation activates AKT and ERK, and promotes resistance to ibrutinib and other agents used in the treatment of Waldenstrom's Macroglobulinemia. *Leukemia*. 2014 Jun 10. doi: 10.1038/leu.2014.187.
103. Cao Y, Hunter ZR, Liu X, et al. CXCR4 WHIM-like frameshift and nonsense mutations promote ibrutinib resistance but do not supplant MYD88 L265P directed signaling in Waldenstrom macroglobulinaemia cells. *Br J Haematol*. 2015 Mar;168(5):701-7.
104. Ghobrial IM, Perez R, Baz R, et al. Phase Ib study of the novel anti-CXCR4 antibody ulocuplumab (BMS-936564) in combination with lenalidomide plus low-dose dexamethasone, or with bortezomib plus dexamethasone in subjects with relapsed or refractory multiple myeloma. *Blood* 2014; 124: 3483 (abstract).
105. Smith EL, Palomba ML, Park J H, et al. A Systemic Xenograft Model of Waldenström's Macroglobulinemia Demonstrates the Potent Anti-Tumor Effect of Second Generation CD19 Directed Chimeric Antigen Receptor Modified T Cells in This Disease. *Proc. American Society for Hematol*. 2014; Abstract 4484.
106. Nagao T, Oshikawa G, Ishida S, et al. A novel MYD88 mutation, L265RPP, in Waldenstrom's macroglobulinemia activates the

NF- $\kappa$ B pathway to upregulate Bcl-xL expression and enhance cell survival. *Blood Cancer J.* 2015; May 15; 5:e314. Doi: 10.1038/bcj.2015.36

## Legends

**Figure 1. Guide to the Primary Therapy of Waldenstrom's Macroglobulinemia.** HV, hyperviscosity; CRYO, Cryoglobulinemia; CAGG, Cold Agglutininemia. BDR (Bortezomib, Dexamethasone, Rituximab); Benda-R (Bendamustine, Rituximab); CaRD (Carfilzomib, Rituximab, Dexamethasone); CDR (Cyclophosphamide, Dexamethasone, Rituximab); FR (Fludarabine, Rituximab), IB (Ibrutinib), R (Rituximab). <sup>1</sup>Consider blood warmers with plasmapheresis in patients with cryoglobulinemia or cold agglutininemia to avoid cryoprecipitation or agglutination. <sup>2</sup>Can be considered in patients who are not wild-type for MYD88<sup>70</sup>, and for those patients without bulky adenopathy or Bing-Neel Syndrome. <sup>3</sup>For patients requiring immediate disease control, consider twice weekly dosed bortezomib for 1-2 cycles, then if patient stable switch to weekly bortezomib to reduce risk of treatment related peripheral neuropathy. In patients not requiring immediate disease control, the use of weekly dosed bortezomib is preferable. Bortezomib should be avoided in patients with disease related neuropathy. Bortezomib should be held for grade  $\geq 2$  treatment-related neuropathy. Acyclovir and famotidine (or equivalent) are strongly recommended for patients on proteasome inhibitor therapy. <sup>4</sup>Rituximab should be held in patients with symptomatic HV, severe CRYO or CAGG, and in asymptomatic patients with serum IgM >4,000 mg/dL to avoid an IgM flare and potentiation of symptoms. Ofatumumab may be considered for rituximab intolerant patients. Consider maintenance rituximab for patients responding to a rituximab-containing regimen. See text for suggested dosing, cycles and scheduling of therapy. A clinical trial should be considered whenever possible.

**Figure 2. Guide to therapy of previously treated Waldenstrom's Macroglobulinemia.** ASCT (Autologous Stem Cell Transplant); RIC (Reduced Intensity Allogeneic Stem Cell Transplant); IB, Ibrutinib; NA; nucleoside-analogue based therapy. <sup>1</sup>Can be considered in patients not previously treated with IB, and who are not wild-type for MYD88, and for those patients without bulky adenopathy or Bing-Neel Syndrome. <sup>2</sup>In patients being considered for an ASCT, stem cell collection should be undertaken before exposure to a nucleoside-analogue. Rituximab should be held in patients with symptomatic HV, severe CRYO or CAGG, and in asymptomatic patients with serum IgM >4,000 mg/dL to avoid an IgM flare and potentiation of symptoms. Ofatumumab may be considered for rituximab intolerant patients. Consider maintenance rituximab for patients responding to a rituximab-containing regimen. Bortezomib should be held for grade  $\geq 2$  treatment-related neuropathy. See text for suggested dosing, cycles and scheduling of therapy. A clinical trial should be considered whenever possible.

**Table 1.** Case presentations of three patients with Waldenstrom Macroglobulinemia.

**Table 2.** Clinical and laboratory findings for 356 newly diagnosed patients meeting consensus diagnostic criteria for WM who presented to the Bing Center for WM clinic (a tertiary care center) at the Dana Farber Cancer Institute, Boston MA. NA; Not Applicable.

**Table 3.** Morbidities mediated by the IgM monoclonal protein in patients


with Waldenstrom's macroglobulinemia.

**Table 4.** A review of systems and considerations for their use in working up patients with Waldenstrom's macroglobulinemia.


**Table 5.** Laboratory and clinical evaluations which can be performed in the initial work-up of all WM patients, and those with particular disease related features.

**Table 6.** A summary of consensus response criteria adopted at the 6<sup>th</sup> International Workshop on Waldenstrom's Macroglobulinemia.<sup>81</sup>

**Figure 1.**



**Figure 2.**



**Table 1.**

**Case 1. WM Patient with IgM-related Peripheral Neuropathy.**

A 67 year-old patient complained of numbness and tingling in his toes that extended above the ankles over several months. Exam revealed diminished pinprick sensation over the soles of his feet. An IgM $\lambda$  monoclonal protein was discovered, with a serum IgM level of 640 mg/dL. A bone marrow (BM) biopsy showed 10% involvement with lymphoplasmacytic lymphoma (LPL), and the MYD88<sup>L265P</sup> mutation was present. Fat-pad biopsy with congo-red staining was negative. An anti-MAG IgM antibody screen was markedly positive with a titer >1:102,400.

### **Case 2. Newly diagnosed WM patient in Hyperviscosity Crisis.**

A 42 year-old male presented with blurry vision and nosebleeds. Exam revealed retinal hemorrhages, adenopathy and splenomegaly. Laboratories revealed a hematocrit of 18%, platelets of 50,000/mm<sup>3</sup>, and leukocyte count of 1500/mm<sup>3</sup>. Serum total protein was high prompting a workup that revealed an IgM $\lambda$  monoclonal protein and serum IgM level of 12,400 mg/dL. CT scans showed bulky adenopathy, and a BM biopsy demonstrated 80% LPL involvement. MYD88<sup>L265P</sup> mutation was present.

### **Case 3. Patient with Refractory WM Disease.**

A 46 year-old male was diagnosed with WM after presentation with fatigue and a hematocrit of 28.6%. BM biopsy showed 90% LPL involvement. CT scans were unremarkable, and serum IgM was 2,780 mg/dL. Treatment with bortezomib, dexamethasone and rituximab was started. After 3 cycles, no treatment response was observed. He then received 2 cycles of cyclophosphamide-based (CDR) therapy without response. Bendamustine was then initiated, and after two cycles serum IgM and hemoglobin levels remained unchanged. Everolimus was then begun and stopped due to non-response and worsening blood counts. A repeat BM biopsy confirmed unchanged WM disease.

**Table 2.**

	<b>Median</b>	<b>Range</b>	<b>Normal Reference Range</b>
Age (yr)	58	32-91	NA
Gender (Male/Female)	215/141		NA
BM involvement	30%	5-95%	NA
Adenopathy	15%		NA
Splenomegaly	10%		NA
IgM (mg/dL)	2,620	270-12,400	40-230
IgG (mg/dL)	674	80-2,770	700-1,600
IgA (mg/dL)	58	6-438	70-400
Serum Viscosity (cp)	2.0	1.1-7.2	1.4-1.9
Hct (%)	35.4%	17.2-45.4%	34.8-43.6%
Plt ( $\times 10^9/L$ )	275	42-675	155-410
Wbc ( $\times 10^9/L$ )	6.4	1.7-22	3.8-9.2
B <sub>2</sub> M (mg/dL)	2.5	0.9-13.7	0-2.7
LDH (U/mL)	313	61-1,701	313-618

**Table 3.**

<b>Properties of IgM Monoclonal Protein</b>	<b>Diagnostic Condition</b>	<b>Clinical Manifestations</b>
Pentameric Structure	Hyperviscosity	Headaches, blurred vision, epistaxis, retinal hemorrhages, leg cramps, impaired mentation, intracranial hemorrhage.
Precipitation on cooling	Cryoglobulinemia (Type I)	Raynaud-like phenomena, acrocyanosis, ulcers, purpura, cold urticaria.
Auto-antibody activity to Myelin Associated Glycoprotein (MAG), Ganglioside M1 (GM1), Sulfatide moieties on peripheral nerve sheaths	Peripheral neuropathies	Sensorimotor neuropathies, painful neuropathies, ataxic gait, bilateral foot drop.
Auto-antibody activity to IgG	Cryoglobulinemia (Type II)	Purpura, arthralgias, renal failure, sensorimotor neuropathies.
Auto-antibody activity to red blood cell antigens	Cold agglutinins	Hemolytic anemia, Raynaud's phenomenon, acrocyanosis, livedo reticularis.
Tissue deposition as amorphous aggregates	Organ Dysfunction	Skin: bullous skin disease, papules, Schnitzler's syndrome.  GI: diarrhea, malabsorption, bleeding.  Kidney: proteinuria, renal failure (light chain component).

Tissue deposition as amyloid fibrils  (light chain component most common)	Organ Dysfunction	Fatigue, weight loss, edema, hepatomegaly, macroglossia, organ dysfunction of involved organs: heart, kidney, liver, peripheral sensory and autonomic nerves.
---	-------------------	---

**Table 4.**

<b>Symptom/Complaint</b>	<b>Implications</b>	<b>Action</b>
Energy level/Changes in activities of daily life.	Anemia, fatigue without anemia	Evaluate for anemia, underlying etiology including iron deficiency, hemolytic anemia (warm and cold antibodies). Consider amyloidosis. Exclude other medical causes of anemia. Hepcidin-related iron deficiency is common in WM. Check iron saturation levels. <sup>32</sup> Patients with low iron saturation levels <10-12% may benefit with parenteral iron infusions. <sup>33</sup>
Constitutional Complaints	Tumor related fever, chills, night sweats.	
Recurrent sinus and bronchial infections.	Chronic sinusitis, usually on the basis of IgA and IgG hypogammaglobulinemia.	Antibiotic support. If patient is refractory to multiple antibiotic courses, required hospitalizations, or infections were life threatening, strongly consider IVIG replacement.

<p>Headaches, blurry vision or visual loss, confusional episodes, epistaxis.</p>	<p>Hyperviscosity</p>	<p>Funduscopy examination for hyperviscosity-related changes, obtain serum IgM and serum viscosity levels.<sup>63,64,118</sup> Serum viscosity levels may be slow to be resulted, not reproducible or lack correlation to serum IgM levels. Consider emergent plasmapheresis for symptomatic hyperviscosity.</p>
<p>Easy bruising, bleeding diathesis.</p>	<p>Thrombocytopenia, Acquired Von Willebrand's Disorder (VWD).</p>	<p>Complete blood count, evaluate for immune thrombocytopenia or hypersplenism if indicated; consider evaluation for VWD; consider amyloidosis.</p>
<p>Progressive symmetrical numbness, tingling, burning, pain feet and hands. Unsteady gait, deficits in motor function.</p>	<p>IgM-related neuropathy or myopathy; amyloidosis.</p>	<p>Obtain anti-MAG, and if negative then anti-GM1, anti-sulfatide IgM antibody studies. Consider obtaining fat pad biopsy and Congo-red stain to evaluate for amyloidosis. Obtain EMG studies and neurology consult.</p>
<p>Raynaud-like symptoms, acrocyanosis, ulcerations on extremities.</p>	<p>Cryoglobulinemia, cold agglutininemia.</p>	<p>Obtain cryoglobulins cold agglutinins. In patients suspected of having cryoglobulins, all studies including quantitative immunoglobulins should be obtained in a warm bath to avoid cryoprecipitation and false lowering of serum IgM levels.</p>

Diarrhea, gastrointestinal cramping.	Malabsorption, secondary to amyloidosis, IgM deposition, tumor involvement. Rarely, autonomic neuropathy on basis of auto-antibody or amyloidosis.	Endoscopy to evaluate small bowel, biopsy to evaluate for amyloidosis, IgM deposition, tumor involvement.
Hearing loss	Hyperviscosity, sensorineural hearing loss, amyloid or tumor deposition, thrombus formation.	Evaluate for anti-hu antibody, MRI to assess for amyloidoma, tumor deposition. Evaluate for hyperviscosity syndrome (as above). Assess for IgM anti-phospholipid antibodies.
Thrombotic events	Antiphospholipid antibody syndrome	Assess for IgM anti-phospholipid antibodies.
Urticaria, papules, dermatitis.	Schnitzler's syndrome (non-pruritic urticaria), IgM or tumor cell infiltration, amyloid deposition.	Skin biopsy, histological examination for tumor cell infiltration, stain for IgM, Congo-red staining for amyloid.

**Table 5.**

<b>Examination</b>	<b>WM Patient Population</b>	<b>Remarks</b>
Beta-2-Microglobulin (B <sub>2</sub> M)	All patients.	Serum B <sub>2</sub> M > 3.0-3.5 mg/L associated with poor prognosis. <sup>87,88</sup> Serum B <sub>2</sub> M is a determinant in WM International Prognostic Staging System. <sup>89</sup> Value of B <sub>2</sub> M in making treatment decisions remains to be clarified.
Blood Urea Nitrogen (BUN), Creatinine	All patients.	If moderate to severe azotemia is present, additional work-up including renal biopsy for deposition of light and heavy chains, amyloid fibrils, and cryoglobulins, and tumor infiltration can be considered.
Bone Marrow Biopsy and Aspiration	All patients.	Morphological evaluation for lymphoplasmacytic cells. Immunohistochemistry and/or flow cytometry demonstrating clonal population. sIgM, CD19, CD20, CD22, and CD79 expression consistent with WM. <sup>90,91</sup> Up to 20% of WM cases may be CD5, CD10, or CD23 positive. <sup>92</sup> MYD88 L265P positive >90% of cases, and can support WM diagnosis.
Cold Agglutinins (CAGG)	Select patients with anemia, reticulocytosis, Raynaud-like symptoms, and/or acrocyanosis.	Present in up to 5% of WM patients.
Complete Blood Counts (CBC)	All patients.	Hb <10 g/dL, and/or PLT count of <100,000/mm <sup>3</sup> related to WM disease can be used to support initiation of therapy based on consensus guidelines. <sup>30</sup>

Cryoglobulins	Select patients with Raynaud-like symptoms, acrocyanosis, peripheral non-healing ulcerations, acrocyanosis, peripheral sensory neuropathy, or erratic serum IgM readings.	Present in 10-15% of WM patients. Patients with cryoglobulinemia should have serum IgM levels obtained on warm bath.
CT Scans of Chest, Abdomen, Pelvis	All patients.	Follow-up CT scans are only necessary for patients with baseline extra-medullary disease or who later are suspected of having extra-medullary disease. Repeat CT scans should be considered in patients with relapse or progression. <sup>31,34</sup>
Cytogenetic Analysis	Select patients with bone lytic lesions, ambiguous bone marrow or laboratory findings concerning for myeloma.	14q32 translocations present in most IgM myeloma cases with t(11;14) being most common, but are absent in WM. <sup>93</sup> MYD88 <sup>L265P</sup> can also be evaluated in BM specimens in ambiguous cases, as it is positive in >90% of WM but absent in IgM myeloma cases. <sup>4,94</sup> 6q deletions common in WM, but prognostic significance and clinical utility is unclear. <sup>95-97</sup>
Electromyography (EMG)	Patients with peripheral sensory neuropathy.	EMG typically shows de-myelinating pattern with anti-myelin IgM-related peripheral sensory neuropathy. Axonal degeneration can be present with amyloid or cryoglobulinemic involvement. <sup>98</sup>
Fat-Pad Biopsy	Select patients with suspected amyloidosis, peripheral neuropathy, nephropathy, unexplained arrhythmias, cardiomegaly or heart failure.	Congo-red staining on fat pad and/or bone marrow biopsy may be used to establish presence of amyloid. Cardiac and renal work-up should be considered if Congo-red staining is positive to clarify extent of amyloid deposition.

Funduscopy Examination	All patients.	Indirect ophthalmoscopy with scleral depression should also be considered in patients with hyperviscosity symptoms, and/or elevated serum IgM levels (>3,000 mg/dL). <sup>99</sup>
Hepatitis testing	Select patients with cryoglobulinemia, and who are being considered for rituximab.	Hepatitis C implicated in some cases of Type II cryoglobulinemia. <sup>100</sup> Hepatitis B testing should be evaluated in all patients who are being considered for rituximab. HBSAg-positive WM patients may be also be candidates for prophylactic anti-viral therapy during rituximab therapy. <sup>101</sup>
Iron, Total Iron Binding Capacity (TIBC)	Select patients with anemia.	Hepcidemia is common in WM and can impact iron absorption and re-utilization. <sup>32</sup> If iron saturation (Iron/TIBC) is <10-12% and not related to GI blood loss may be candidate for parenteral iron therapy. <sup>33</sup>
Liver Function Tests (LFTs)	All patients.	
Lumbar puncture with CSF Examination.	Select patients with suspected Bing-Neel Syndrome.	Brain MRI with gadolinium should first be done to evaluate for parenchymal and meningeal involvement, and evaluate risk for herniation. Cytological examination, white cell count with differential, flow cytometry, PCR for IgH rearrangements, as well as evaluation for MYD88 <sup>L265P</sup> can be obtained to identify WM cells in CSF. <sup>102</sup> If CSF positive, and Brain MRI is negative, MRI of entire spinal axis can be considered to evaluate for tumor presence if brain MRI is negative.

MYD88 Genotyping		MYD88 L265P positive >90% of cases by AS-PCR, and can support WM diagnosis. <sup>4-8</sup> MYD88 <sup>WT</sup> is associated with inferior overall survival; and lower response rates and progression free survival to ibrutinib in WM patients. <sup>14,70</sup> Sanger sequencing for non-L265P MYD88 mutations can be considered in WM patients who by AS-PCR are negative for MYD88 <sup>L265P</sup> as other types of MYD88 mutations can occur in WM. <sup>70,106</sup>
Positron Em/ission Tomography (PET) Scan	Select patients.	No role for routine PET scans in WM unless disease transformation is suspected. <sup>31,34</sup>
Serum IgM, IgA, IgG levels (Quantitative)	All patients.	Obtain on warm bath for patients suspected of having cryoglobulinemia. Serum IgM levels can flare following rituximab, and should be carefully monitored in WM patients with high serum IgM levels (>4,000 mg/dL). <sup>31,34</sup> Quantitative serum IgM levels can be used as surrogate marker for disease burden assessment. <sup>84</sup> Serum IgA, IgG levels are subnormal in most WM patients and impacted by certain treatments. <sup>80</sup> Patients with severely depressed IgG and recurring sinobronchial infections may benefit with intravenous immunoglobulin (IVIG) infusions.
Reticulocyte Count (RC)	Select patients with anemia.	For patients with high reticulocyte counts, consider hemolysis work-up. Extravascular hemolysis is common in WM, and LDH, and haptoglobin levels may be normal. Consider cold agglutinin testing, and if negative obtain direct and indirect Coombs evaluation. Splenic capture may also be etiological for reticulocytosis in WM patients with hypersplenism.

Serum Free Light Chains (SFLC)	Select patients with light-chain nephropathy, amyloidosis.	May be useful for serial assessment of treatment impact in patients with amyloidosis or light chain deposition disease.
Serum Myelin Associated Glycoprotein (MAG) titers	Select patients with peripheral sensory neuropathy, some forms of ataxia.	Positive in 25-50% of WM patients with disease related peripheral neuropathy. <sup>69,103</sup> If negative, may consider further testing for anti-ganglioside M1 (GM1) and anti-sulfatide IgM antibodies.
Serum Protein Electrophoresis (SPEP)	All patients.	Demonstration of serum monoclonal IgM protein required for diagnosis of WM. Monoclonal IgM protein may be used as surrogate marker for assessing disease burden. <sup>84</sup> Migration pattern of serum IgM and formation of higher order complexes on gels can complicate serial IgM protein assessment in some patients. <sup>85</sup> Absence of serum IgM monoclonal protein by SPEP and immunofixation studies following therapy can be used to support attainment of a complete response (see Table 6). <sup>81</sup>
Serum Viscosity	Select patients with suspected hyperviscosity symptoms and findings.	May be useful in assessing patients with hyperviscosity symptoms, though slow to be resulted and erratic readings commonly occur. <sup>63,64</sup> Serum viscosity levels may be slow to be resulted, not reproducible or lack correlation to serum IgM levels. Serum IgM levels are more expedient and reliable for assessing patients with suspected hyperviscosity syndrome. Funduscopic examination should be done in all patients with hyperviscosity syndrome. <sup>99</sup>

**Table 6.**

<b>Complete Response</b>	<b>CR</b>	IgM in normal range, and disappearance of monoclonal protein by immunofixation; no histological evidence of bone marrow involvement, and resolution of any adenopathy / organomegaly (if present at baseline), along with no signs or symptoms attributable to WM. Reconfirmation of the CR status is required by repeat immunofixation studies.
<b>Very Good Partial Response</b>	<b>VGPR</b>	A $\geq$ 90% reduction of serum IgM and decrease in adenopathy/organomegaly (if present at baseline) on physical examination or on CT scan. No new symptoms or signs of active disease.
<b>Partial Response</b>	<b>PR</b>	A $\geq$ 50% reduction of serum IgM and decrease in adenopathy/organomegaly (if present at baseline) on physical examination or on CT scan. No new symptoms or signs of active disease.
<b>Minor Response</b>	<b>MR</b>	A $\geq$ 25% but < 50% reduction of serum IgM. No new symptoms or signs of active disease.
<b>Stable Disease</b>	<b>SD</b>	A <25% reduction and <25% increase of serum IgM without progression of adenopathy/organomegaly, cytopenias or clinically significant symptoms due to disease and/or signs of WM.
<b>Progressive Disease</b>	<b>PD</b>	A $\geq$ 25% increase in serum IgM by protein confirmed by a second measurement or progression of clinically significant findings due to disease (i.e. anemia, thrombocytopenia, leukopenia, bulky adenopathy/organomegaly) or symptoms (unexplained recurrent fever $\geq$ 38.4°C, drenching night sweats, $\geq$ 10% body weight loss, or hyperviscosity, neuropathy, symptomatic cryoglobulinemia or amyloidosis) attributable to WM.



# blood

Prepublished online May 22, 2015;  
doi:10.1182/blood-2015-01-553974

## How I treat Waldenstrom macroglobulinemia

Steven P. Treon

---

Information about reproducing this article in parts or in its entirety may be found online at:  
[http://www.bloodjournal.org/site/misc/rights.xhtml#repub\\_requests](http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests)

Information about ordering reprints may be found online at:  
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:  
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>

---

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include digital object identifier (DOIs) and date of initial publication.